11.4: Bernoulli Trials and Binomial Probability Distributions

Bernoulli trials:

A *Bernoulli trial* is an experiment with exactly two possible outcomes. We refer to one of the outcomes as a *success* (S) and to the other as a *failure* (F). We'll call the probability of success p and the probability of failure q. In other words,

P(S) = p and P(F) = 1 - p = q. (Note that S and F are complements of one another.)

Example 1: Roll a single die. Consider rolling a 5 to be a success.

Example 2: Roll a pair of dice. Consider success to be rolling a sum of 7, 11, or 12.

Consider what happens when a Bernoulli trial is repeated several times. Now we can discuss the probability of a particular number of successes.

Suppose we repeat a Bernoulli trial 6 times and each time the trial is independent of the others. From the multiplication principle, what is the probability of getting SFSSSF?

What is the probability of getting SSSFFS?

What is the probability of having 4 successes in the 6 trials?

Example 3: If a single die is rolled 6 times, what is the probability of getting four 5's?

Binomial experiments:

Now let's write all this down in a more precise way:

A sequence of experiments is called a sequence of Bernoulli trials, or a *binomial experiment*, if

- 1. The same experiment is repeated a fixed number of times. (Each repetition is called a trial.)
- 2. Only two outcomes are possible on each trial.
- 3. The probability of success *p* for each trial is a constant.
- 4. All trials are independent.

Probabilities in Bernoulli Trials:

If the probability of success is p and the probability of failure is q, then the probability of exactly x successes in p Bernoulli trials is

$$P(x \text{ successes}) = C_{n,x} p^x q^{n-x}$$
.

The binomial formula:

For any natural number n,

$$(a+b)^{n} = C_{n,0}a^{n}b^{0} + C_{n,1}a^{n-1}b^{1} + C_{n,2}a^{n-2}b^{2} + \dots + C_{n,n}a^{0}b^{n}$$
$$= \sum_{i=0}^{n} C_{n,i}a^{n-i}b^{i}$$

Notice the similarity between this formula and the formula for probabilities in a sequence of Bernoulli trials. This is why such a sequence is called a *binomial experiment*.

Example 4: If a single fair die is rolled 10 times, what is the probability of

- a. Exactly three 4's?
- b. At most three 4's?
- c. At least one 4?

Example 5: In the United States, about 9% of people have blood type B+. If 20 people donate blood, what is the probability that exactly three of them are B+? At least three?

The binomial distribution:

We can now generalize Bernoulli trials and determine probability distributions.

In a binomial experiment with n trials and probability of success p, we can create a binomial distribution table, with the variable x representing the number of successes.

Example 6: Suppose a fair die is rolled three times and a success is considered to be rolling a number divisible by 3.

Expected value of the binomial distribution:

We can determine the average number of successes (or the *expected value* of x) for any binomial distribution.

Expected Value of the Binomial Distribution:

In a binomial experiment with n trials and probability of success p, the expected value, or mean, is

$$E(x) = \mu = np$$
.

Standard Deviation of the Binomial Distribution:

$$\sigma = \sqrt{npq} .$$

Example 7: Roll a single fair die 10 times. Consider rolling a 4 to be a success. Find the expected number of successes.

