


### 3.3: Increasing and Decreasing Functions and the First Derivative Test


#### Increasing and decreasing functions:

A function  $f$  is said to be *increasing* on the interval  $(a,b)$  if, for any two numbers  $x_1$  and  $x_2$  in  $(a,b)$ ,  $f(x_1) < f(x_2)$  whenever  $x_1 < x_2$ . A function  $f$  is *increasing at  $c$*  if there is an interval around  $c$  on which  $f$  is increasing.



all tangent lines have positive slopes

A function  $f$  is said to be *decreasing* on the interval  $(a,b)$  if, for any two numbers  $x_1$  and  $x_2$  in  $(a,b)$ ,  $f(x_1) > f(x_2)$  whenever  $x_1 < x_2$ . A function  $f$  is *decreasing at  $c$*  if there is an interval around  $c$  on which  $f$  is decreasing.



all tangent lines have negative slopes


Notice that wherever a function is increasing, the tangent lines have positive slope.  
Notice that wherever a function is decreasing, the tangent lines have negative slope.

This means that we can use the derivative to determine the intervals where a function is increasing and decreasing.

Increasing/Decreasing Test: Let  $f$  be a function that is continuous on the closed interval  $[a,b]$  and differentiable on the open interval  $(a,b)$ .

- If  $f'(x) > 0$  for every  $x$  in  $(a,b)$ , then  $f$  is increasing on  $(a,b)$ .
- If  $f'(x) < 0$  for every  $x$  in  $(a,b)$ , then  $f$  is decreasing on  $(a,b)$ .
- If  $f'(x) = 0$  for every  $x$  in  $(a,b)$ , then  $f$  is constant on  $(a,b)$ .

**Example 1:**  $f(x) = x^2$



$$f'(x) = 2x$$

$$2x > 0 \text{ for } x > 0$$

$$2x < 0 \text{ for } x < 0$$

From graph,  
 $f$  is decreasing on  $(-\infty, 0)$   
 and increasing on  $(0, \infty)$

#### Steps for Determining Increasing/Decreasing Intervals

1. Find all the values of  $x$  where  $f'(x) = 0$  or where  $f'(x)$  is not defined. Use these values to split the number line into intervals.
2. Choose a test number  $c$  in each interval and determine the sign of  $f'(c)$ .
  - If  $f'(c) > 0$ , then  $f$  is increasing on that interval.
  - If  $f'(c) < 0$ , then  $f$  is decreasing on that interval.

Note: Three types of numbers can appear on your number line:

- 1) Numbers where the function is defined and the derivative is 0. (These are critical numbers.)
- 2) Numbers where the function is defined and the derivative is undefined. (These are also critical numbers.)
- 3) Numbers where the function is undefined. (These are NOT critical numbers.)

#### **First derivative test:**

This procedure determines the relative extrema of a function  $f$ .

#### First derivative test:

Suppose that  $c$  is a critical number of a function  $f$  that is continuous on an open interval containing  $c$ .

- If  $f'(x)$  changes from positive to negative across  $c$ , then  $f$  has a relative maximum at  $c$ .
- If  $f'(x)$  changes from negative to positive across  $c$ , then  $f$  has a relative minimum at  $c$ .
- If  $f'(x)$  does not change sign across  $c$ , then  $f$  does not have a relative extreme at  $c$ .

From number line, using 1<sup>st</sup> derivative test,  $f$  has a relative max at  $-6$  and a relative min at  $2$ . Find the  $y$ -values:  $f(-6) = (-6)^3 + 6(-6)^2 - 36(-6) + 18 = 234$   
 $f(2) = 2^3 + 6(2)^2 - 36(2) + 18 = -22$  3.3.3


**Example 2:** Determine the intervals on which  $f(x) = x^3 + 6x^2 - 36x + 18$  is increasing and decreasing. Find the relative extrema.

Domain:  $(-\infty, \infty)$

$$\begin{aligned} f'(x) &= 3x^2 + 12x - 36 \\ &= 3(x^2 + 4x - 12) \\ &= 3(x+6)(x-2) \end{aligned}$$

Set  $f'(x) = 0$ :  $0 = 3(x+6)(x-2)$   
 $x = -6, 2$

Critical numbers:  $-6, 2$



$(-\infty, -6)$ : Test number:  $x = -7$

$$\begin{aligned} f'(-7) &= 3(-7)^2 + 12(-7) - 36 \\ &= 3(49) - 84 - 36 = 147 - 120 \\ &= 27 (+) \end{aligned}$$

$$\begin{aligned} f'(-7) &= 3(-7+6)(-7-2) \\ &= 3(-1)(-9) = 27 (+) \\ &= (+)(-)(-) \end{aligned}$$

$(-6, 2)$ : Test number:  $x = 0$

$$f'(0) = 3(0)^2 + 12(0) - 36 = -36 (-)$$

$(2, \infty)$ : Test number:  $x = 3$


$$\begin{aligned} f'(3) &= 3(3+6)(3-2) \\ &= 3(9)(1) = 27 (+) \end{aligned}$$

$f$  is increasing on  $(-\infty, -6)$  and on  $(2, \infty)$ .  
 $f$  is decreasing on  $(-6, 2)$ .

**Example 3:** Determine the intervals on which  $g(x) = x^3 - 6x^2 + 12x - 8$  is increasing and decreasing. Find the relative extrema. Domain:  $(-\infty, \infty)$

$$\begin{aligned} g'(x) &= 3x^2 - 12x + 12 \\ &= 3(x^2 - 4x + 4) \\ &= 3(x-2)^2 \end{aligned}$$

Critical number:  $2$



$(-\infty, 2)$ : Test number  $x = 0$ :

$$g'(0) = 3(0-2)^2 = 3(4) = 12 (+)$$

$(2, \infty)$ : Test number:  $x = 3$

$$g'(3) = 3(3-2)^2 = 3(1)^2 = 3 (+)$$

$g$  is increasing on  $(-\infty, \infty)$

No relative extrema

**Example 4:**

Determine the intervals on which  $g(x) = x^{2/5}$  is increasing and decreasing. Find the relative extrema.

$$g(x) = x^{2/5} = \sqrt[5]{x^2}$$

Domain:  $(-\infty, \infty)$

$$g'(x) = \frac{2}{5} x^{-3/5} = \frac{2}{5\sqrt[5]{x^3}}$$

where is  $g'$  undefined? at  $x=0$

Where is  $g'$  zero? nowhere

Critical value: 0

$(-\infty, 0)$ : Test  $x = -1$

$$g'(-1) = \frac{2}{5\sqrt[5]{(-1)^3}} = \frac{2}{5(-1)} = -\frac{2}{5} \quad (-)$$

$(0, \infty)$ : Test  $x = 1$

$$g'(1) = \frac{2}{5\sqrt[5]{1^3}} = \frac{2}{5} \quad (+)$$



$g$  is increasing on  $(0, \infty)$ .  
 $g$  is decreasing on  $(-\infty, 0)$ .

Relative min at  $x=0$ .

Find  $y$ -value:

$$g(0) = \sqrt[5]{0^2} = 0$$

Relative minimum:  $g(0) = 0$

**Example 5:**

Determine the intervals on which  $f(x) = x + \frac{4}{x}$  is increasing and decreasing. Find the relative extrema.

**Example 6:** Find the local extremes of  $g(x) = (x^2 - 4)^{\frac{2}{3}}$ . Where is it increasing and decreasing?

**Example 7:** Find the relative extremes of  $f(x) = \frac{1}{2}x - \sin x$  on the interval  $(0, 2\pi)$ . Where is it increasing and decreasing on that interval?