## 3.5: Descriptive Measures for Populations; Use of Samples (also z-scores and measures of position)

## Population mean, population variance, and population standard deviation:

Recall:

Population mean:

$$\mu = \frac{\sum_{i=1}^{n} x_{i}}{n} = \frac{x_{1} + x_{2} + \ldots + x_{n}}{n}$$

$$\chi_{i} = mu$$

Sample mean:

$$\overline{x} = \frac{\sum_{i=1}^{n} x_i}{n} = \frac{x_1 + x_2 + \dots + x_n}{n}$$

Population variance:

$$\sigma^{2} = \frac{\sum_{i=1}^{n} (x_{i} - \mu)^{2}}{n}$$

Sample variance:

$$s^{2} = \frac{\sum_{i=1}^{n} (x_{i} - \overline{x})^{2}}{n-1}$$

Population standard deviation:

$$\sigma = \sqrt{\frac{\sum_{i=1}^{n} (x_i - \mu)^2}{n}}$$

$$\sigma = \sqrt{\frac{\sum_{i=1}^{n} (x_i - \mu)^2}{n}}$$

Sample standard deviation:

$$s = \sqrt{\frac{\sum_{i=1}^{n} (x_i - \overline{x})^2}{n-1}}$$

## Parameters and statistics:

Definition:

A parameter is a descriptive measure for a population.

A *statistic* is a descriptive measure for a sample.

Parameters: population mean u, population varione oz, population std-der Statistics: sample mean x, sample varione 12, sample sta. deviction a

## The z-score:

The *z*-score for a data point represents the number of standard deviations that lie between the data point and the mean. The *z*-score is sometimes known as the standardized value, and it allows us to compare data points from different distributions.


You can think of the *z*-score as representing the "distance from the mean," with distance measured in standard deviations. A positive *z*-score indicates the data point lies above the mean; a negative *z*-score indicates the data point lies below the mean.

Therefore, for bell-shaped distributions (from Empirical Rule),

- about 68% of the data points have z-scores between –1 and 1;
- 96% about  $\frac{1}{2}$  of the data points have z-scores between -2 and 2;
- 99.7% about 2 of the data points have z-scores between -3 and 3.

For all distributions (from Chebyshev's Rule),

- at least 75% of the data points have z-scores between –2 and 2;
- at least 88.9% of the data points have z-scores between -3 and 3.



Note: The z-score is unitless. All distributions of z-scores have mean 0 and standard deviation 1.

Example 1: Suppose a data set has mean 52 and standard deviation 8. Find the z-scores for the scores 44, 64, 38, and 52.  $M = 52 \times 9 = 9$ 

scores 44, 64, 38, and 32.

For 
$$x = 4A$$
:  $Z = \frac{x - M}{\sigma} = \frac{44 - 52}{9} = \frac{12}{9} = \frac{12}{9}$ 

For  $x = 6A$ :  $Z = \frac{x - M}{\sigma} = \frac{(a4 - 52)}{9} = \frac{12}{9} = \frac{12}{9} = \frac{12}{9}$ 

For  $x = 39 - 52$ 

For  $x = 39 - 52$ 

For  $x = 52$ ,  $z = \frac{52 - 52}{9} = \frac{0}{9} = \frac{0}{9}$ 

**Example 2:** In 2014, the mean of the ACT mathematics test was 20.9 and the standard deviation was 5.3. In the same year, the mean of the SAT mathematics test was 513 and the standard deviation was 120. Suppose Tamara, a high school student, received a score of 24 on the ACT mathematics test, and 600 on the SAT mathematics test. On which test did she perform better?

(ACT data from the National Center for Education Statistics, <a href="https://nces.ed.gov/programs/digest/d14/tables/dt14\_226.50.asp?current=yes">https://nces.ed.gov/programs/digest/d14/tables/dt14\_226.50.asp?current=yes</a>; SAT data from the College Board, <a href="https://www.collegeboard.org/program-results/2014/sat">https://www.collegeboard.org/program-results/2014/sat</a>)

We calculate her z-score on each test:

ACT her score: x = 24  $\mu = 20.9$   $\sigma = 5.3$   $Z = \frac{x-\mu}{\sigma} = \frac{24-20.9}{5.3} \approx 0.586$ SAT her score: x = 600  $\mu = 513$   $\sigma = 120$   $Z = \frac{x-\mu}{\sigma} = \frac{600-513}{120} = 0.735$ Relative to the reticued energes, she scored batter on the SAT (because her z-score is higher)