

6.3: Working with Normally Distributed Variables

Recall: The z-score of a data point is its distance from the mean, measured in standard deviations.




Standardizing the values of a normal distribution:

In a normal distribution with mean μ and standard deviation σ , where x is a data value, the z-score is

$$z = \frac{x - \mu}{\sigma}$$

The area under a normal curve between $x = a$ and $x = b$ is the same as the area under the standard normal curve between the z-score for a and the z-score for b .

Example 1: Consider a normal curve with mean 7 and standard deviation 2.



Properties of Normal Probability Distributions:

1. $P(a \leq x \leq b)$ = area under the curve from a to b .
2. $P(-\infty \leq x \leq \infty) = 1$ = total area under the curve.
3. $P(x = c) = 0$.

Note: $P(a \leq x \leq b) = P(a \leq x < b) = P(a < x \leq b) = P(a < x < b)$


Example 2: Dusty Dog Food Company ships dog food to its distributors in bags whose weights are normally distributed with a mean weight of 50 pounds and standard deviation 0.5 pound. If a bag of dog food is selected at random from a shipment, what is the probability that it weighs

- a) More than 51 pounds?
- b) Less than 49 pounds?
- c) Between 49 and 51 pounds?
- d) What is the percentage of dog food bags that weigh more than 51 pounds?

$$\mu = 50$$

$$\sigma = 0.5$$

a)



Find z-score for $x = 51$: $z = \frac{x - \mu}{\sigma} = \frac{51 - 50}{0.5} = 2.00$


Look up $z = 2.00$ in table
 $\Rightarrow A_1 = 0.4772$

$$P(X > 51) = P(Z > 2.00)$$

$$= 0.5 - A_1 = 0.5 - 0.4772$$

$$= \boxed{0.0228}$$

b)




Could calculate z-score

$$\text{For } x = 49, \quad z = \frac{x - \mu}{\sigma} = \frac{49 - 50}{0.5} = -2.00$$

or use symmetry:

$$P(X < 49) = \boxed{0.0228} \quad \text{Same as part (a)}$$

c) Between 49 and 51 lbs:



Because of symmetry, $A_1 = A_2$

From Part (a), $A_1 = 0.4772$

$$P(49 < X < 51) = 2(0.4772)$$

$$= \boxed{0.9544}$$

Example 3: The medical records of infants delivered at a certain hospital show that the infants' birth weights in pounds are normally distributed with a mean of 7.4 and a standard deviation of 1.2.

- a) What percentage of infants at this hospital weighed more than 9.2 pounds at birth?
- b) What percentage of infants at this hospital weighed less than 8 pounds at birth?
- c) What percentage of infants at this hospital weighed between 8 and 10 pounds at birth?

Important: The z -score is the number of standard deviations between the data point and the mean.

Example 4: A variable is normally distributed with mean 83 and standard deviation 24.

- a) Find and interpret the quartiles.
- b) Find and interpret the 98th percentile.
- c) Find and interpret the first and second deciles.
- d) Find the value that 72% of all possible values of the variable exceed.
- e) Find two values of the variable that divide the area into a middle area of 0.90 and two outside areas of 0.05 each.

Example 5: The GPA of the senior class of a certain high school is normally distributed with a mean of 2.7 and a standard deviation of 0.4 point. If a senior in the top 10% of his or her class is eligible for admission to any state university, what is the minimum GPA that a senior should have to ensure eligibility to a state university?