



8.4.1

8.4: Bayes' Formula

Previously, when we discussed conditional probabilities, we considered the probability of a "later" event (in a probability tree), given that an "earlier" event occurred.

Now, we will consider the probability of an event "earlier" in the tree, given that a "later" event occurs.

Example 1: A box contains 4 white marbles and 3 red marbles. Two marbles are drawn in succession without replacement. What is the probability of drawing a red marble on the first draw given that a red marble is drawn on the second draw?

$$P(R_1 | R_2) = \frac{P(R_1 \cap R_2)}{P(R_2)} = \frac{\frac{3}{6} \cdot \frac{2}{5}}{\frac{4}{6} \cdot \frac{2}{5} + \frac{3}{6} \cdot \frac{2}{5}} = \frac{\frac{6}{30}}{\frac{12}{30} + \frac{6}{30}} = \frac{\frac{6}{30}}{\frac{18}{30}} = \frac{6}{18} = \frac{1}{3}$$


↓ "prob of the intersection divided by the prob. of the given"

We've just used something a theorem known as *Bayes' Theorem*. You don't need to memorize it, as it is just the conditional probability formula $P(A|B) = \frac{P(A \cap B)}{P(B)}$.

Bayes' Formula:

$$P(A|B) = \frac{\text{product of probabilities on path to B through A}}{\text{sum of all branch products leading to B}} = \frac{P(A \cap B)}{P(B)}$$

Example 2: The picture tubes for the Pulsar 19-inch television sets are manufactured in three locations and then shipped to the main plant for final assembly. Plants A, B, and C supply 50%, 30%, and 20%, respectively, of the picture tubes. The quality-control department of the company has determined that 1% of the picture tubes produced by plant A are defective, whereas 2% of the tubes produced by plants B and C are defective. If a Pulsar 19-inch color television is selected at random and the picture tube is found to be defective, what is the probability that the picture tube was manufactured in plant C?



$$P(C|D) = \frac{P(C \cap D)}{P(D)}$$

$$= \frac{0.20(0.02)}{0.50(0.01) + 0.30(0.02) + 0.20(0.02)}$$

$$= \frac{0.004}{0.015}$$

$$\approx \boxed{0.267}$$