

3.6: Polynomial and Rational Inequalities

Polynomial and rational inequalities are examples of *nonlinear* inequalities. They cannot be solved simply by adding, subtracting, multiplying, or dividing both sides by the same quantity.

Examples: $\frac{x-5}{x+3} \geq 0$ or $x^2 - 7x + 12 > 0$

Can we do this?

$$\begin{aligned} x^2 - 7x + 12 &> 0 \\ (x-3)(x-4) &> 0 \\ x-3 &> 0 \text{ or } x-4 > 0 \\ x &> 3 \text{ or } x > 4 \end{aligned}$$

Does this give us the right answer?

If not, where did our logic go wrong?

To solve a polynomial or rational inequality:

- Rearrange so that one side is 0.
- If the nonzero side involves quotients, write it with a common denominator.
- If possible, factor the nonzero side. If it's a quotient, factor the numerator and factor the denominator.
- Find all the numbers that make the expression zero or undefined.
- Use these values to divide the number line into intervals.
- In each interval, choose a test number to determine whether the expression is positive or negative. It's easiest to use the factored form.
- Use this information to make a "sign chart".
- Use the sign chart to determine what values make the original inequality true.
- Write the solution in interval notation.

Example 1: Solve $9 - x^2 \leq 0$.

Example 2: Solve $x^2 - 7x > -12$.


$$x^2 - 7x + 12 > 0$$

To solve this, graph $y = x^2 - 7x + 12$

$$y = (x-3)(x-4)$$

x-intercepts: 3, 4

opens up.



Solution Set: $(-\infty, 3) \cup (4, \infty)$

Solve $x^2 - 7x + 12 \leq 0$


Sol'n Set: $[3, 4]$

Example 3: Solve $-2(x+5)(x-1)(x-4)^2 \geq 0$

Bonus
on Final

$$f(x) = -2(x+5)(x-1)(x-4)^2$$

Lead term: $-2(x)(x)(x)^2 = -2x^4$



Solution Set:
 $[-5, 1] \cup \{4\}$

Example 4: Solve $3x^2 - 7x < 6$.

Example 5: Solve $\frac{2x+6}{x-2} \geq 0$.

Example 6: Solve $\frac{3+x}{3-x} \geq 1$.

Example 7: Solve $\frac{1}{x-2} < \frac{2}{x+2}$.