
Making Predictions about the Population Mean from the Sample Mean:



population mean: μ

population standard deviation: σ


We can definitely use the sample mean as an estimate for the population mean, but how accurate will it be?

If the sample size is large(n > 30) or the population is normal, then the sample mean, \overline{x} , is normal(approximately) with mean equal to the population mean and standard deviation of $\frac{\sigma}{\sqrt{n}}$, called the <u>Standard Error of the Mean</u>.

We want to come up with an interval centered at the sample mean, \bar{x} , that contains the population mean, μ , with a certain probability. This interval is called a confidence interval for the population mean.

The confidence interval can be written as $\bar{x} \pm E$ or $(\bar{x} - E, \bar{x} + E)$, and the probability that it contains the population mean is called its confidence level, and it's usually written as a $(1-\alpha)$ confidence interval. The width of the confidence interval is $2 \cdot E$.

So we want $P(\overline{x}-E<\mu<\overline{x}+E)=1-\alpha$, and this is equivalent to $P(\mu-E<\overline{x}<\mu+E)=1-\alpha$. Now let's subtract the mean and divide by the standard deviation. $P\left(-\frac{E}{\frac{\sigma}{E}}<\frac{\overline{x}-\mu}{\frac{\sigma}{E}}<\frac{E}{\frac{\sigma}{E}}\right)=1-\alpha$ or just $P\left(-\frac{E}{\frac{\sigma}{E}}< Z<\frac{E}{\frac{\sigma}{E}}\right)=1-\alpha$.



 $z_{\frac{\omega}{2}}$ is called a critical value so that $P(Z>z_{\frac{\omega}{2}})=\frac{\alpha}{2}$, and therefore that $P(-z_{\frac{\omega}{2}}< Z< z_{\frac{\omega}{2}})=1-\alpha$.

So for our confidence interval to work, it must be that $\frac{E}{\frac{\sigma}{\sqrt{n}}} = z_{\frac{\pi}{2}}$, or $E = z_{\frac{\pi}{2}} \cdot \frac{\sigma}{\sqrt{n}}$. So the

$$(1-\alpha)$$
 confidence interval can be written as $\overline{x} \pm z_{\frac{\alpha}{2}} \cdot \frac{\sigma}{\sqrt{n}}$ or $(\overline{x} - z_{\frac{\alpha}{2}} \cdot \frac{\sigma}{\sqrt{n}}, \overline{x} + z_{\frac{\alpha}{2}} \cdot \frac{\sigma}{\sqrt{n}})$.

For a 90% confidence interval, $1-\alpha=.9$, so $\alpha=.1$, and $\frac{\alpha}{2}=.05$. $z_{.05}=1.645$.

For a 95% confidence interval, $1-\alpha = .95$, so $\alpha = .05$, and $\frac{\alpha}{2} = .025$. $z_{.025} = 1.96$.

For a 99% confidence interval, $1-\alpha = .99$, so $\alpha = .01$, and $\frac{\alpha}{2} = .005$. $z_{.005} = 2.576$.





Examples:

1. A sample of size 16 is randomly selected from a normal population with a standard deviation of 5.23. If the sample mean is 15.1, then construct confidence intervals with the following confidence levels.

a) 90%


width of the confidence interval:

b) 95%

width of the confidence interval:

c) 99%

width of the confidence interval:



Finding the Sample Size for a Desired Level of Accuracy:

Suppose that you would like to know how large a sample is required to estimate a population mean with an error of at most E at a $(1-\alpha)$ confidence.

$$z_{\mathcal{L}} \cdot \frac{\sigma}{\sqrt{n}} \leq E$$

$$\frac{1}{\sqrt{n}} \leq \frac{E}{z_{\text{e/s}} \cdot \sigma}$$

$$\sqrt{n} \ge \frac{z_{\%} \cdot \sigma}{E}$$

$$n \ge \left(\frac{z_{\%} \cdot \sigma}{E}\right)^2$$



" I got the instructions from my Statistics Professor. He was 80% confident that the true location of the restaurant was in this neighborhood."

Examples:

1. You would like to estimate the population mean of a population with a standard deviation of 1.2 with an error of at most .05 with 90% confidence. How large of a sample should you take?

$$n \ge \left(\frac{z_{4} \cdot \sigma}{E}\right)^2$$

TOYOU.

2. You would like to estimate the population mean of a population with a standard deviation of 1.8 with an error of at most .08 with 95% confidence. How large of a sample should you take?

$$n \ge \left(\frac{z_{\frac{\alpha}{2}} \cdot \sigma}{E}\right)^2$$