3.1. Radian Measure

Thursday, February 14, 2019 8:02 AM

1°



Divide the circle into 360 equal parts.

1 radian



Circle, radius R = 2 inches

↳ length of arc = 2 inches

central angle has measure 1 radian







Radius = R, arc length = s

θ = angle intercepted by this arc.

The measure of θ in radians is: $\theta = \frac{s}{R}$



$$\theta = \frac{s}{R} = \frac{2\pi \cancel{R}}{\cancel{R}} = 2\pi$$

So, $360^\circ = 2\pi$ radians

$\longrightarrow 180^\circ = \pi$ radians

$1^\circ = \frac{\pi}{180}$ radians

1 radian $= \frac{180}{\pi}$ degrees

To convert from

degree $\xrightarrow{\text{multiply by } \frac{\pi}{180}}$ radian

radian $\xrightarrow{\text{multiply by } \frac{180}{\pi}}$ degree

Ex. Convert the following to radians

(a) 30° $\frac{\pi}{6}$

(b) 45° $\frac{\pi}{4}$

(c) 60° $\frac{\pi}{3}$

(d) 90° $\frac{\pi}{2}$

(e) 120° $\frac{2\pi}{3}$

(f) 135° $\frac{3\pi}{4}$

Ex. Convert the following to degrees

(a) $\frac{5\pi}{6} \cdot \frac{180}{\pi}$ 150°

(b) $\frac{7\pi}{6}$ 210°

(c) $\frac{11\pi}{6}$ 330°

(d) $\frac{3\pi}{4}$ 135°

(e) $\frac{5\pi}{4}$ 225°

(f) $\frac{7\pi}{4}$ 315°



HW: Fill out the rest of the circle.

Import family of angles that have the same reference angle.

$\frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$. Reference angle: $\frac{\pi}{4}$

$\frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3}$ _____: $\frac{\pi}{3}$

$\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ _____: $\frac{\pi}{6}$

θ	$\sin\theta$	$\cos\theta$	$\tan\theta$	$\csc\theta$	$\sec\theta$	$\cot\theta$
$\frac{\pi}{6} = 30^\circ$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	2	$\frac{2\sqrt{3}}{3}$	$\sqrt{3}$
$\frac{\pi}{4} = 45^\circ$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	$\sqrt{2}$	$\sqrt{2}$	1
$\frac{\pi}{3} = 60^\circ$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{2\sqrt{3}}{3}$	2	$\frac{\sqrt{3}}{3}$

E.g. Find the exact value of:

(a) $\tan\frac{2\pi}{3}$

Reference angle: $\frac{\pi}{3}$ → $\tan\frac{\pi}{3} = \sqrt{3}$

Quadrant: II → $\tan\frac{2\pi}{3} = \boxed{-\sqrt{3}}$

(b) $\sec\frac{7\pi}{4}$ Reference angle: $\frac{\pi}{4}$ → $\sec\frac{\pi}{4} = \sqrt{2}$

Quadrant: IV → $\sec\frac{7\pi}{4} = \sqrt{2}$.

(c) $\sin\left(-\frac{7\pi}{6}\right)$

Coterminal: $-\frac{7\pi}{6} + 2\pi = \frac{5\pi}{6}$

Reference angle: $\frac{\pi}{6}$ → $\sin\frac{\pi}{6} = \frac{1}{2}$

Quadrant: II → $\sin\left(-\frac{7\pi}{6}\right) = \frac{1}{2}$.

(d) $\cos\left(-\frac{14\pi}{3}\right)$

Coterminal: $-\frac{14\pi}{3} + 3\cdot 2\pi = -\frac{14\pi}{3} + 6\pi = \frac{4\pi}{3}$

Reference angle: $\frac{\pi}{3}$ → $\cos\frac{\pi}{3} = \frac{1}{2}$

Quadrant: III → $\cos\left(-\frac{14\pi}{3}\right) = -\frac{1}{2}$

(e) $\cos(3\pi) = \cos(\pi) = -1$

$\cos(2019\pi) = \cos(\pi) = -1$

$\cos(2020\pi) = \cos(0) = 1$

$\cos(k\pi)$ $\begin{cases} 1 : k \text{ is even} \\ -1 : k \text{ is odd} \end{cases}$

$\sin(k\pi) = 0$

Formula for finding reference angle of angle θ in $[0, 2\pi]$



$\theta' = \theta$



$\theta' = \pi - \theta$



$\theta' = \theta - \pi$



$\theta' = 2\pi - \theta$